

Fapas[®] – Food Microbiology

INSTRUCTIONS FOR THE PREPARATION OF SAMPLES

Beef, Chicken, Meat, Fish, Herbs (parsley), Egg, Salad, Rice and Ready To Eat (RTE) Meal

These samples require a **rehydration** stage before you start the analysis.

TREAT EACH SAMPLE AS A WHOLE – DO NOT SUB-SAMPLE and ensure you rinse the container thoroughly.

For each sample for an **ENUMERATION test**: add 10 ml (+/- 0.2 ml) buffered peptone water.

Important Note: this gives you a sample which is equivalent to 10 g of a routine sample.

For each sample for a **DETECTION test (except tests for *Vibrio parahaemolyticus*)**: add 20 ml (+/- 0.2 ml) buffered peptone water.

Important Note: this gives you a sample which is equivalent to 25 g of a routine sample.

For each sample for a **DETECTION test for *Vibrio parahaemolyticus***: add 20 ml (+/- 0.2 ml) **ASPW** (Alkaline Saline Peptone Water) or equivalent

Important Note: this gives you a sample which is equivalent to 25 g of a routine sample

Then:

- Gently invert the sample a few times to aid rehydration.
- Leave the sample to stand at room temperature for 30 minutes (+/- 2minutes).

The sample is now ready to test using your usual procedure.

Milk Powder and Animal Feed Test Materials

These samples require a **resuscitation** stage before you start the analysis.

TREAT EACH SAMPLE AS A WHOLE – DO NOT SUB-SAMPLE and ensure you rinse the container thoroughly.

Add the sample to the blender / homogeniser bag.

For each sample for an **ENUMERATION test**: add 90 ml (+/- 2 ml) of your usual diluent, rinsing the sample container with part of the diluent.

Important Note: this makes a 1/10 dilution

For each sample for a **DETECTION test**: add 225 ml (+/- 5 ml) of your usual pre-enrichment / enrichment broth, rinsing the sample container with part of the broth.

Important Note: this makes a 1/10 dilution.

Then:

- Leave the sample to stand at room temperature for 30 minutes (+/- 2 minutes).

The sample is now ready to test using your usual procedure.

Ground Pepper, Chocolate, Chocolate Powder, Infant Formula and Flour

These samples can be analysed without any special preparation.

TREAT EACH SAMPLE AS A WHOLE – DO NOT SUB-SAMPLE and ensure you rinse the container thoroughly.

For each sample for an **ENUMERATION test** OR a **DETECTION test**.

The sample is ready to test using your usual procedure.

Sponge Swab for Detection

TREAT EACH SAMPLE AS A WHOLE – DO NOT SUB-SAMPLE.

Add 10 ml (+/- 0.2 ml) buffered peptone water directly to the sponge in the container.

Then:

- Leave the sample to stand at room temperature for 30 minutes (+/- 2 minutes).

The sample is now ready to test using your usual procedure.

Cotton Swab for APC (Aerobic Plate Count)

TREAT EACH SAMPLE AS A WHOLE – DO NOT SUB-SAMPLE.

Add 10 ml (+/- 0.2 ml) buffered peptone water to the sample and vortex thoroughly for 30 seconds.

Then:

- Leave the sample to stand at room temperature for 30 minutes (+/- 2 minutes) to rehydrate.
- Vortex the sample for 10 seconds.

The sample is now ready to test using your usual procedure.

Fruit Juice

TREAT EACH SAMPLE AS A WHOLE – DO NOT SUB-SAMPLE and ensure you rinse the vial thoroughly.

Add 1ml (from a 10ml (+/- 0.2 ml) aliquot) of buffered peptone water to the sample vial.

Then:

- Leave the sample to stand at room temperature for 1 minute (+/- 10 seconds)
- Carefully rinse the vial contents twice with a pastette (Pasteur pipette)
- Transfer the full contents from the vial into the 10ml buffered peptone water (from which the initial 1ml amount was taken) to give a final volume of 10ml (+/- 0.2 ml)

The sample is now ready to test using your usual procedure.

Cheese (for *E. coli* 0157 tests)

These samples require a **rehydration** stage before you start the analysis.

TREAT EACH SAMPLE AS A WHOLE – DO NOT SUB-SAMPLE and ensure you rinse the container thoroughly.

For each sample for a **DETECTION test** add 20 ml (+/- 0.2 ml) buffered peptone water.

Important Note: this gives you a sample which is equivalent to 25 g of a routine sample.

Then:

- Gently invert the sample a few times to aid rehydration.
- Leave the sample to stand at room temperature for 30 minutes (+/- 2minutes).

The sample is now ready to test using your usual procedure.

Soft Cheese (for *Listeria monocytogenes* tests)

These samples can be analysed without any special preparation.

TREAT EACH SAMPLE AS A WHOLE – DO NOT SUB-SAMPLE and ensure you rinse the container thoroughly.

For each sample for an **ENUMERATION test** OR a **DETECTION test**.

The sample is ready to test using your usual procedure.